

El-Shahawy Y*

Department of Obstetrics and Gynecology, Ain Shams University, Abbasyia, Cairo, Egypt

Dates: Received: 29 February, 2016; Accepted: 05 March, 2016; Published: 08 March, 2016

*Corresponding author: Yasser El-Shahawy, lecturer of Obstetrics and Gynecology, Faculty of Medicine, Ain Shams University, Abbasyia, Cairo, Egypt

www.peertechz.com

Keywords: Endometrial volume; VOCAL; Pregnancy rate; IVF; ICSI

Research Article

Reproducibility of Endometrial Volume by VOCAL in the Prediction of Success Rate of IVF/ICSI in Women with Unexplained Infertility

Abstract

Objective: To explore the role of estimating endometrial volume by Virtual Organ Computer-aided Analysis (VOCAL) and its correlation with endometrial thickness, on the day of hCG, in prediction of IVF/ICSI success.

Patients and Methods: It was a prospective study that was carried out at Al-Amin Maternity Hospital, Saudi Arabia. Endometrial thickness and volume were measured in women undergoing an IVF/ICSI cycle, on the day of HCG, using the 3D transvaginal ultrasound probe. Women were divided as regards to endometrial volume calculated into 3 subgroups; < 3ml, 3-5ml, and >5ml, also according to endometrial thickness into 3 groups; <7mm, 7-12mm and >12mm. Success rate of IVF/ICSI was compared between all groups.

Results: The current study included 150 consented women with no significant difference in clinic-demographic characteristics between all subgroups. The pregnancy rates between the three groups of endometrial volume; <3ml, 3-5ml, and >5 ml was; 28.6%, 29.9% and 25% respectively. Pregnancy rates between the three groups of endometrial thickness was 18.2%, 25.5% and 43.3% for <7mm, 7-12mm and >12mm respectively. There was no significant difference in pregnancy rates between all groups in relation to both endometrial volume and endometrial thickness.

Conclusions: Endometrial thickness on day of hCG is a better predictor of success rate of IVF/ICSI cycles than endometrial volume.

Introduction

There are many causes of infertility. In the majority of couples, an apparent male or a female factor explaining their failure to conceive is usually present, in 10–25% there will be no explanation. Within this cohort of women there is a spectrum of diseases ranging from those women with a decreased fertility that will get pregnant with time to those in whom pregnancy is very difficult with available medical practice [1].

Pregnancy in In Vitro Fertilization (IVF) / intracytoplasmic sperm injection (ICSI) programs needs embryos of good quality and a receptive endometrium [2]. Despite the outstanding progress in these techniques, the implantation rate per embryo is constant at very low at 20 % [3]. While the evaluation of quality of embryos and correct grading is performed in the lab before transfer, the evaluation of uterine lining receptivity is somewhat difficult. Transvaginal sonogram (TVS) is an ideal non-invasive technique, but with deficient specificity [2-4]. The utilization of 3 D ultrasonography to test the details of the uterine cavity surpasses the diagnostic ability of Two Dimensional (2D) ultrasound [5,6]. There is a rising trend to utilize noninvasive methods as TVS, to assess the endometrial thickness, diagnose dysfunctional uterine bleeding (DUB), adenomyosis, endometrial polyps & fibroids [7]. Another important potential of 3D TVS is volume calculation utilizing the Virtual Organ Computer-aided Analysis (VOCAL) even in irregularly shaped organs. This

technique has been shown to be more precise than 2D-volume estimation [5].

Many studies have stated a high level of reproducibility and accuracy of endometrial volume measurement using 3D ultrasound [8,9]. The aim of this study was to evaluate endometrial volume measured by VOCAL, on the day of human chorionic gonadotrophin (hCG), to endometrial thickness in the prediction of IVF/ICSI success.

Patients and Methods

This was a prospective study performed at Al-Amin Maternity Hospital, Saudi Arabia and included 150 women who were presented for having unexplained infertility and were scheduled for IVF/ICSI done using ejaculate sperms. This study is performed over a 2-year period, between Jan 2013 and Jan 2015. Endometrial thickness and volume were measured in women undergoing an IVF/ICSI cycle, on the day of HCG, using the 3D transvaginal ultrasound probe. Women were divided as regards to endometrial volume calculated into 3 subgroups; < 3ml, 3-5ml, and >5ml. Also according to the thickness of the endometrium they were divided into 3 groups; <7mm, 7-12mm and >12mm. Success rate of IVF/ICSI was compared between all groups.

Inclusion criteria

1. Women under 40 year old.

2. Women with unexplained infertility.

Exclusion criteria

1. loss to follow-up.
2. Any patient with uterine abnormality was excluded (e.g. fibroid, mullerian anomaly).
3. Patients with a previous poor response.
4. Those with difficult embryo transfer.
5. Women with a known systemic disease.
6. Women with more than 3 failed IVF.

Informed consent was taken from every women, after they were fully discussed about the nature, scope the potential risks of this clinical trial.

Ovulation induction protocol used was the long luteal phase agonist protocol. Women received Decapeptyl 0.1 mg/day starting on day 20 of the cycle till the day of hCG intake. After down regulation, daily 150- 300 IU of human menopausal gonadotrophin (hMG) was commenced for 7 days, then the needed dose was given according to the follicular response. 10.000 units of hCG (Pregnyl; Nile CO., Cairo Egypt) were administered I.M. when more than two follicles reached 17mm in mean diameter. Ovum pickup (OPU) using transvaginal ultrasound was performed 36 hour after hCG intake [10]. Embryo transfer was performed on day 5 after OPU using the Wallace catheter (H.G.Wallace Ltd, West Sussex, UK). Daily progesterone injection (100 mg, progesterone; Steris, Phoenix, AZ, USA) was given for luteal phase support. A serum B-hCG test was performed to check pregnancy two weeks after embryo transfer. Clinical pregnancy was diagnosed 3 weeks after a positive test by the presence of a gestational sac with fetal echoes and pulsations on ultrasound.

Endometrial evaluation on the day of hCG intake

The patients were checked using the TV 3D probe, the endometrial thickness was assessed at the maximum thickness between the highly reflective interfaces of the endometrial-myometrial junction [8]. Three-dimensional volumetry was done with an empty bladder, the patient was examined at the Special Care Center of The Fetus by the same sonographer in the lithotomy position using Voluson Pro 720 GENERAL ELECTRIC USA, transducer with frequency range 5-8 MHZ. Sonar is done to study uterine size, shape and exclusion of any uterine or ovarian pathology. The 3D image was obtained by switching on the 3D volume mode and defining the region of interest by a movable sector on the screen. This sector has the shape of a truncated cone which can be manipulated to ensure that the whole of the endometrial cavity was included in the volume sampling while the patient remain still and the probe is held stationary. Volume sampling lasted about 4 seconds, during that time the conventional 2D plane was rotated through 180 with the rotation axis oriented exactly along the longitudinal axis of the vaginal probe. The data was stored digitally on the internal disc drive for subsequent analysis after the ultrasound probe is removed. For the purpose of volume calculation, 3D data was retrieved and presented in multi-planer display mode which simultaneously displays 3 perpendicular planes on the screen. The actual volume was calculated by the built-in computer program using VOCAL. This is a rotational method based on rotation in given steps (6,9,15,30) on a given orthogonal plane (A,B OR C). The endometrial

volume was measured in plane A by delineating the endometrial margin at the endometrial-myometrial interface from the fundus to the internal cervical is in a number of parallel slices which are 1-2 mm apart [8].

The hospital ethics committee approved the study. All patients gave their informative consent before entering into the study.

Statistical analysis

Retrieved data were recorded on an investigative report form. The data were analyzed with SPSS[®] for Windows[®], version 15.0 (SPSS, Inc, USA). Significance level was set at 0.05.

Results

The current study was conducted on 150 women presented at Al-Amin Maternity Hospital, Saudi Arabia during the period between Jan 2013 and Jan 2015. The mean age was 29.3 ± 3.4 (range 20-40), the mean duration of infertility was 7.3 ± 1.9 years. The number of embryos transferred was 2. The overall fertilization rate was 71.5%, clinical pregnancy rate; 31.8%. Patients were divided into three subgroups according to endometrial volume calculated; <3ml, 3-5ml, >5ml and according to endometrial thickness into three subgroups; <7mm, 7-12mm, >12mm. The mean age, period of infertility and number of embryos transferred in all subgroups is shown in Table 1 and there was no significant difference in these parameters Table 2 shows the number of pregnancies and pregnancy rates in each subgroup and differences were found to be non-significant. Table 3 shows a non-significant difference in the mean endometrial volume

Table 1: Patient characteristics in the 3 subgroups of endometrial volume & thickness.

Endometrial thickness	Women age	Infertility duration
<7 mm	28.6 ± 2.8	7.5 ± 2.3
7-12 mm	26.9 ± 2.1	8.9 ± 2.5
>12 mm	31.1 ± 1.4	6.9 ± 3.1
Endometrial volume		
<3ml	27.6 ± 3.8	7.7 ± 2.4
3-5 ml	30.3 ± 4.1	6.3 ± 2.1
>5 ml	28.5 ± 2.7	8.7 ± 1.8

Table 2: Pregnancy rate in the 3 subgroups of endometrial volume and endometrial thickness.

Endometrial thickness	Number	Pregnant	Non pregnant	Ratio %
<7 mm	22	4	18	18.2
7-12 mm	98	25	73	25.5
>12 mm	30	13	17	43.3
Endometrial volume				
<3ml	35	10	25	28.6
3-5 ml	87	26	61	29.9
>5 ml	28	7	21	25

Table 3: Comparison between endometrial volume and endometrial thickness in the pregnant and non-pregnant cases.

	Pregnant	Non-pregnant	P-value
Endometrial thickness	11.8	9.8	<0.005 Significant
Endometrial volume	4.8	4.2	>0.005 NS

* Analysis using independent student's t-test. NS = non-significant.

but significant mean endometrial thickness in the pregnant as compared to the non-pregnant group.

Discussion

It is well known that the quality of embryos and the receptivity of the endometrium determine the success rate of IVF/ ICSI and successful implantation [11]. In the last few years, there has been a rising interest in evaluation of endometrial receptivity by methods of ultrasound technology. Many researchers tried to assess endometrial receptivity through many sonographic parameters, as endometrial thickness, but with low specificity [12]. Utilizing 2D transvaginal ultrasound to evaluate endometrial thickness does not comprise the total endometrial volume [2]. Other investigators stated a 12.6% as an error rate of for two dimensional volume measurements, whereas the error rate for three dimensional volume measurement was 6.4% only [13]. In addition the benefit of reproducibility of 3D endometrial volume measurement has been established by many researchers [8,9]. Many reports examined the volume of the endometrium on the days of down regulation, ovum retrieval and embryo transfer. In the current study there was no statistically significant difference in endometrial volume in women who conceived as compared to those who did not, but there was a significant difference in endometrial thickness between the two groups. This is in partial agreement with previous studies that reported no predictive value of measuring endometrial volume and thickness on IVF success [14,15]. When subdividing the women into three groups according to endometrial volume; <3ml, 3-5ml, >5 ml, also according to the thickness of the endometrium <7mm, 7- 12mm, >12mm, no statistically significant difference in pregnancy rates was found as regards to endometrial volume but there was a significant difference as regards to endometrial thickness which doesn't match with other investigators who subdivided the women similarly [16]. In the current study when analyzing endometrial volume & thickness assessment of women with grade I embryos, endometrial thickness was found to be better than endometrial volume in predicting pregnancy. Hence the validity of endometrial volume for canceling ET or cryopreserving embryos is questionable. Other authors reported that endometrial volume and thickness significantly higher in women who conceived as compared to those who did not conceive. However in agreement with our results, it was reported no significant reduction in pregnancy rates with endometrial volume < 2ml when compared to > 2ml [17]. In conclusion, 3D endometrial volume assessment is an easy, practical and reproducible method of endometrial evaluation in IVF programs. However endometrial volume is not a better predictor of IVF success when compared to endometrial thickness measured on day of hCG.

References

- Hull MG, Glazener CM, Kelly NJ, Conway DI, Foster PA, et al. (1985) Population study of causes, treatment and outcome of infertility. *Br Med J (Clin Res Ed)* 291: 1693-1697.
- Friedler S, Schenker JG, Herman A, Lewin A (1996) The role of

ultrasonography in the evaluation of endometrial receptivity following assisted reproductive treatments: a critical review. *Hum Reprod Update* 2: 323-335.

- Pope CS, Cook EK, Army M, Novak A, Grow DR (2004) Influence of embryo transfer depth on in vitro fertilization and embryo transfer outcomes. *Fertil Steril* 81: 51-58.
- Sundstrom P (1998) Establishment of a successful pregnancy following in-vitro fertilization with an endometrial thickness of no more than 4 mm. *Hum Reprod* 13: 1550-1552.
- Jurkovic D, Geipel A, Gruboeck K, Jauniaux E, Natucci M, et al. (1995) Three-dimensional ultrasound for the assessment of uterine anatomy and detection of congenital anomalies: a comparison with hysterosalpingography and two-dimensional sonography. *Ultrasound Obstet Gynecol* 5: 233-237.
- Lev-Toaff AS, Pinheiro LW, Bega G, Kurtz AB, Goldberg BB (2001) Three-dimensional multiplanar sonohysterography: comparison with conventional twodimensional sonohysterography and X-ray hysterosalpingography. *J Ultrasound Med* 20: 295-306.
- Alcazar JL, Galvan R (2009) Three-dimensional power Doppler ultrasound scanning for the prediction of endometrial cancer in women with postmenopausal bleeding and thickened endometrium. *Am J Obstet Gynecol* 200: 44.e-1-6.
- Lee A, Sator M, Kratochwil A, Deutinger J, Vytiska-Binsdorfer E, et al. (1997) Endometrial volume change during spontaneous menstrual cycles: volumetry by transvaginal three-dimensional ultrasound. *Fertil Steril* 68: 831-835.
- Yaman C, Ebner T, Sommergruber M, Polz W, Tews G (2000) Role of three-dimensional ultrasonographic measurement of endometrium volume as a predictor of pregnancy outcome in an IVF-ET program: a preliminary study. *Fertil Steril* 74: 797-801.
- Mansour RT, Aboulghar MA, Serour GI, Fahmi I, Ramzy AM, et al. (1996) Intracytoplasmic sperm injection using microsurgically retrieved epididymal and testicular sperm. *Fertil Steril* 65: 566-572.
- Ozturk O, Bhattacharya S, Saridogan E, Jauniaux E, Templeton A (2004) Role of utero ovarian vascular impedance: predictor of ongoing pregnancy in an IVF-embryo transfer program. *Reprod Biomed Online* 9: 299-305.
- Remohi J, Ardiles G, Garcia-Velasco JA, Gaitan P, Simon C, et al. (1997) Endometrial thickness and serum oestradiol concentrations as predictors of outcome in oocyte donation. *Hum Reprod* 12: 2271-2276.
- Riccabona M, Nelson TR, Pretorius DH (1996) Three-dimensional ultrasound: accuracy of distance and volume measurements. *Ultrasound Obstet Gynecol* Jun 7: 429-434.
- Schild RL, Indefrei D, Eschweiler S, Van der Ven H, Fimmers R, et al. (1999) Three-dimensional endometrial volume calculation and pregnancy rate in an in-vitro fertilization programme. *Hum Reprod* 14: 1255-1258.
- Schild RL, Knobloch C, Dorn C, Fimmers R, van der Ven H, et al. (2001) Endometrial receptivity in an in vitro fertilization program as assessed by spiral artery blood flow, endometrial thickness, endometrial volume, and uterine artery blood flow. *Fertil Steril* 75: 361-366.
- Raga F, Bonilla-Musoles F, Casan EM, Klein O, Bonilla F (1999) Assessment of endometrial volume by three-dimensional ultrasound prior to embryo transfer: clues to endometrial receptivity. *Hum Reprod* 14: 2851-2854.
- Zollner U, Zollner KP, Specketer MT, Blissing S, Muller T, et al. (2003) Endometrial volume as assessed by threedimensional ultrasound is a predictor of pregnancy outcome after in vitro fertilization and embryo transfer. *Fertil Steril* 80: 1515-1517.

Copyright: ©2016 El-Shahawy Y. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: El-Shahawy Y (2016) Reproducibility of Endometrial Volume by VOCAL in the Prediction of Success Rate of IVF/ICSI in Women with Unexplained Infertility. *J Gynecol Res Obstet* 2(1): 014-016.