2023 | Volume Volume - 8 - Issue Issue - 3

In this issue

Case Report

Open Access Case Report PTZAID:ACG-8-214

Gelatinous peritoneal disease secondary to appendiceal mucinous adenocarcinoma: a case study and review of the literature

Published On: November 10, 2022 | Pages: 055 - 058

Author(s): Chérihane Dassouli*, Fatimaezzahra Aboutarik, Adil Ait Errami, Sofia Oubaha, Zouhour Samlani and Khadija Krati

Pseudomyxoma peritonei also referred to as gelatinous ascites, is a rare disorder, described for the first time by R. Wyerth in 1884. It is characterized by the presence of mucous disseminated throughout the peritoneal cavity generally arising from the rupture of an appendicular mucocele. Pseudomyxoma peritonei can be asymptomatic, discovered during a laparotomy. The ...

Abstract View Full Article View DOI: 10.17352/2455-2283.000114

Open Access Case Report PTZAID:ACG-8-213

Severe polycystic liver disease: An unsual cause of chronic Budd-Chiari Syndrome

Published On: October 29, 2022 | Pages: 052 - 054

Author(s): Aboutarik Fatima Ezzahra*, Dassouli Cherihane, Ait Errami Adil, Oubaha Sofia, Samlani Zouhour and Krati Khadija

Autosomal dominant polycystic kidney disease is a hereditary disease, characterized by the development of cysts in the renal parenchyma with extra-renal manifestations. Liver damage is rarely a source of complications. A Budd-Chiari syndrome could occur following the compression of the supra-hepatic veins by the cysts. It is an exceptional mechanical complication. W ...

Abstract View Full Article View DOI: 10.17352/2455-2283.000113

Open Access Case Report PTZAID:ACG-8-212

Perianal paget’s disease: A case report

Published On: September 28, 2022 | Pages: 050 - 051

Author(s): Pilar Navajas Hernández, Teresa Valdés Delgado, Jesús Machuca Aguado, Ricardo González-Cámpora and Federico Argüelles Arias*

The incidence of extramammary Paget's disease (EMPD) is very low. It is very important to distinguish between primary Paget's disease and secondary process. An 85-year-old man was consulted for the presence of an erythematous plaque located in the anal and gluteal area, confirming Paget cells in the biopsy. ...

Abstract View Full Article View DOI: 10.17352/2455-2283.000112

Case Study

Open Access Case Study PTZAID:ACG-8-211

Risk factors associated with response to highly active antiretroviral therapy after 24 months of administration to HIV, HIV/HBV and HCV patients in Kumba Health District, South West Region of Cameroon

Published On: September 26, 2022 | Pages: 037 - 049

Author(s): Adamu Ndongho Ndifontiayong*, Innocent Mbulli Ali, Jean Baptiste Sokoudjou, Karimo Ousenu, Jerimiah Ndimumeh Mbogwe, Ousenu Karimo and Christopher Bonglavnyuy Tume

Hepatitis B (HBV) and C (HCV) are two other forms of infections for which co-infection in HIV has been associated with alteration of the immune response, increased risk of progression to liver diseases, and increased risk of hepatotoxicity associated with antiretroviral therapy. This study aimed to establish the prevalence of hepatitis B surface antigen (HBsAg) and h ...

Abstract View Full Article View DOI: 10.17352/2455-2283.000111