

In this issue

Research Article

[Open Access](#) [Research Article](#) PTZAID:ACN-8-162

Studying the efficiency of carbon nano enterosorbents in the model of experimental renal failure

Published On: December 28, 2022 | Pages: 036 - 040

Author(s): NT Abilaikhanova*, GA Tusupbekova, A Beissova, B Ussipbek, AM Mukhitdinov, A Yessenbekova, ZA Balmaganbet, B Ursheeva, GK Atanbayeva and ZB Yessimsiitova

The current article examines the effectiveness of nanoenterosorbent to correct violations of the functional state of the kidneys in the experimental renal failure model. The obtained data open up opportunities for further research aimed at studying the possibility of using nanoenterosorbent in practical medicine as a new nanoenterosorbent and a means of drug delivery. ...

[Abstract View](#) [Full Article View](#) [DOI: 10.17352/acn.000062](#)

Review Article

[Open Access](#) [Review Article](#) PTZAID:ACN-8-159

Cellular interorganelle crosstalk in health and disease states: A glimpse on nephrology-related conditions

Published On: May 13, 2022 | Pages: 007 - 027

Author(s): Fateme Shamekhi Amiri*

Cellular interorganelle crosstalk in medical sciences is a new discussion of mechanisms and pathways of physiological functions and pathogenesis of diseases. The organelles ("mitochondria", nucleus, lysosome, endoplasmic reticulum, Golgi apparatus) are members of such functional units that are needed to perform specific tasks. Mitochondria are essential metabolic orga ...

[Abstract View](#) [Full Article View](#) [DOI: 10.17352/acn.000059](#)

[Open Access](#) [Case Report](#) PTZAID:ACN-8-160

Syndrome of Inappropriate Antidiuretic Hormone secretion due to hydroxyurea

Published On: September 05, 2022 | Pages: 028 - 030

Author(s): Hakan Ozer*, Yasin Ozturk, Ismail Baloglu and Kultigin Turkmen

Hyponatremia is the most common electrolyte disorder in clinical practice: Syndrome of Inappropriate Antidiuretic Hormone Secretion (SIADH) is the most common cause of hyponatremia, especially in euvoletic patients. Drugs are among the most common causes of SIADH. While our patient was using hydroxyurea due to polycythemia vera, he was diagnosed with SIADH due to hydr ...

[Abstract View](#) [Full Article View](#) [DOI: 10.17352/acn.000060](#)

[Open Access](#) [Observational Study](#) PTZAID:ACN-8-161

The occurrence and influencing factors of fatigue and sleep disturbance in maintenance hemodialysis patients

Published On: September 17, 2022 | Pages: 031 - 035

Author(s): Dong Kaixuan*, Chen Xi, Zhou Rong and Yu Xiaoyong

Objective: To investigate the occurrence and influencing factors of fatigue and sleep disturbance in Maintenance Hemodialysis (MHD) patients. Methods: A total of 170 patients with end-stage renal disease who underwent MHD treatment in the hemodialysis room of Shaanxi Provincial Hospital of Traditional Chinese Medicine from October 2021 to March 2022 were selected as ...

[Abstract View](#) [Full Article View](#) [DOI: 10.17352/acn.000061](#)

[Open Access](#) [Observational Study](#) PTZAID:ACN-8-158

Acute kidney infarction: Not so rare renal disease. A single-center experience with endovascular fibrinolytic therapy

Published On: February 17, 2022 | Pages: 001 - 006

Author(s): Scarpioni R*, De Amicis S, Bodini FC, Albertazzi V and Michieletti E

Renal Infarction (RI), a rare cause of renal damage characterized by the abrupt interruption of flow in the renal artery, is often recognized late or may even remain undiagnosed since symptoms are non-specific and may be confused with other pathologies, such as pyelonephritis or nephrolithiasis. In situ thrombosis and thromboembolism are the main causes, but often the ...

[Abstract View](#)

[Full Article View](#)

[DOI: 10.17352/acn.000058](#)