

In this issue

Research Article

[Open Access](#) [Research Article](#) PTZAID:IJDCR-12-158

## **Cholera: Determinants of the Disease and Prospects for Control in the Congo-Ubangui River Corridor in the Republic of Congo**

Published On: June 02, 2026 | Pages: 017 - 027

Author(s): Medard Amona\*, Yolande Voumbo Matoumona Mavoungou, Tatia Adoua Doukaga, Martin Ekat, Albert Mabilia, Michel Mankou, Hama Nemet Ondzotto, Armel Itoua, Gilius Axel Aloumba, Guy Roger Kinouani, Riha Lena Okamba Afissofo, Stevie Loundou, Albert Boungou, Jean Médard Nkankou, Guy Abel Bongo, Roland Bienvenu OSSIBI IBARA and Pascal Ibata

A major public health threat along the Congo-Ubangui river corridor, cholera is exacerbated by inadequate sanitation infrastructure and high population mobility. Our study analyzes the determinants of the disease in these riverside areas with a view to developing integrated control strategies. A descriptive and analytical cross-sectional study was conducted from Marc ...

[Abstract View](#) | [Full Article View](#) | [DOI: 10.17352/ijdcr.000058](#)

[Open Access](#) [Research Article](#) PTZAID:IJDCR-12-157

## **Stevens-Johnson Syndrome and Toxic Epidermal Necrolysis Reactions to Ibuprofen and Acetaminophen: Clinical, Laboratory Assessment Using In Vitro Lymphocyte Toxicity Assay to Personalized Therapy**

Published On: May 01, 2026 | Pages: 001 - 016

Author(s): Neuman Manuela\*, Salisbury Roger and Misha Montfort

Introduction: Stevens-Johnson Syndrome (SJS) and Toxic Epidermal Necrolysis (TEN) are serious cutaneous adverse reactions (SCARs). SJS is defined when 10% of the body's skin is involved. When epidermal detachment involves 30% or more, it is called TEN. The syndrome affects 1-12 cases per million patients. The risk of death for SJS is 5-10%, and for TEN, 25% -50%. We u ...

[Abstract View](#) | [Full Article View](#) | [DOI: 10.17352/ijdcr.000057](#)